arXiv:1211.1224v1 [math.AT] 6 Nov 2012

Combinatorial stratifications and minimality of 2-arrangements

Karim Adiprasito *
Inst. Mathematics, FU Berlin
adiprasito@math.fu-berlin.de

October 25, 2012

Abstract

We prove that the complement of any affine 2-arrangement in $\mathbb{R}^d$ is *minimal*, that is, it is homotopy equivalent to a cell complex with as many i-cells as its i-th Betti number. To this end, we prove that the Björner–Ziegler complement complexes, induced by combinatorial stratifications of any essential 2-arrangement, admit perfect discrete Morse functions. Our results extend previous work by Falk, Dimca–Papadima, Hattori, Randell, and Salvetti–Settepanella, among others.

1 Introduction

In this paper, we study the complement $\mathscr{A}^{\mathrm{c}} := \mathbb{R}^d \setminus \mathscr{A}$ of any 2-arrangement $\mathscr{A}$ in $\mathbb{R}^d$. A c-arrangement is a family of distinct affine subspaces of $\mathbb{R}^d$, all of codimension c, with the property that the codimension of the non-empty intersection of any subset of $\mathscr{A}$ is a multiple of c. For example, after identifying $\mathbb{C}$ with $\mathbb{R}^2$, any family of hyperplanes in $\mathbb{C}^d$ can be viewed as a 2-arrangement in $\mathbb{R}^{2d}$. However, not all 2-arrangements arise this way, cf. [GM88, Sec. III, 5.2], [Zie93].

Subspace arrangements and their complements have been extensively studied in several areas of mathematics. Thanks to the work by Goresky and MacPherson [GM88], the homology of $\mathscr{A}^{\mathrm{c}}$ is well understood; it is determined by the *intersection poset* of the arrangement, which is the set of all nonempty intersections of its elements, ordered by reverse inclusion. In fact, the intersection poset determines even the homotopy type of the compactification of $\mathscr{A}$ [ZŽ93]. It does, however, not determine the homotopy type of the complement of $\mathscr{A}^{\mathrm{c}}$, even if we restrict ourselves to complex hyperplane arrangements [Ryb11]. Understanding the homotopy type of $\mathscr{A}^{\mathrm{c}}$ is thus a challenging goal.

A standard approach to study the homotopy type of a topological space X is to find a *model* for it, that is, a CW complex homotopy equivalent to it. By cellular homology any model of a space X must use at least $\beta_i(X)$ i-cells for each i, where β_i is the i-th Betti number. A natural question arises: Is the complement of an arrangement *minimal*, that is, does it have a model with *exactly* $\beta_i(X)$ i-cells for all i?

Building on previous work by Hattori [Hat75], Falk [Fal93] and Cohen–Suciu [CS98], around 2000 Dimca–Papadima [DP03] and Randell [Ran02] independently showed that the complement of any complex hyperplane arrangement is a minimal space. Roughly speaking, the idea is to consider the distance to a complex hyperplane in general position as a Morse function on the Milnor fiber to establish a Lefschetz-type theorem. An elegant inductive argument then completes their proof.

On the other hand, the complement of an arbitrary subspace arrangement is, in general, *not* minimal. In fact, complements of subspace arrangements might have arbitrary torsion in cohomology (compare [GM88, Sec. III, Thm. A]). Moreover, complements for 2-arrangements *of pseudospheres* are not minimal either, cf. Björner–Ziegler [BZ92]. This naturally leads to the following question:

Problem I (Minimality)**.** *Is the complement $\mathscr{A}^{\mathrm{c}}$ of an arbitrary c-arrangement $\mathscr{A}$ minimal?*

The interesting case is $c = 2$. In fact, if c is not 2, the complements of c-arrangements, and even c-arrangements of pseudospheres, are easily shown to be minimal; see Section 7.

---

*Supported by the DFG within the research training group “Methods for Discrete Structures” (GRK1408) and by the Romanian NASR, CNCS – UEFISCDI, project PN-II-ID-PCE-2011-3-0533.

In 2007, Salvetti–Settepanella [SS07] proposed a combinatorial approach to Problem I, based on Forman's discretization of Morse theory [For98]. Discrete Morse functions are defined on regular CW complexes, rather than on manifolds; instead of critical points, they have combinatorially-defined *critical faces*. Any discrete Morse function with c_i critical i-faces on a complex C yields a model for C (the *Morse complex*) with exactly c_i i-cells [For98, Cor. 3.5]. Salvetti–Settepanella studied discrete Morse functions on the *Salvetti complexes* [Sal87], which are models for complements of complexified real arrangements. Remarkably, they found that all Salvetti complexes admit *perfect* discrete Morse functions, that is, functions with exactly $\beta_i(\mathscr{A}^{\mathrm{c}})$ critical i-faces. The associated Morse complexes now yield minimal models.

Unfortunately, this tactic does not extend to the generality of complex hyperplane arrangements. However, models for complex arrangements, and even for c-arrangements, have been introduced and studied by Björner and Ziegler [BZ92]. These models boil down to the Salvetti complex in the complexified-real case. While our notion of the combinatorial stratification is slightly more restrictive than Björner–Ziegler's, cf. Section 2.1, it still includes most of the combinatorial stratifications studied in [BZ92]. For example, we still recover the $\mathbf{s}^{(1)}$-stratification which gives rise to the Salvetti complex in the complexified-real case. With these tools in our hand, we can tackle Problem I combinatorially:

Problem II (Optimality of classic models)**.** *Are there perfect discrete Morse functions on the Björner–Ziegler models for the complements of arbitrary c-arrangements?*

We are motivated by the recent discovery that discrete Morse theory, together with the possibility of taking (an arbitrary number of) barycentric subdivisions, is a powerful tool to determine homology [AB11, AB12a], even on stratified spaces. On the other hand, there are several difficulties to overcome. In fact, Problem II is more ambitious than Problem I in many respects:

- Few CW complexes, even among those having trivial topology, admit perfect discrete Morse functions. For example, many 3-balls [Goo68] and many contractible 2-complexes [Zee64] are not collapsible.
- There are few results in literature predicting the existence of perfect Morse functions. For example, it is not known whether any subdivision of the 4-simplex is collapsible [Kir95, Pr. 5.5].
- Solving Problem II could help in obtaining a more explicit picture of the attaching maps for the minimal model; compare Salvetti–Settepanella [SS07] and Yoshinaga [Yos07].

In the present paper, we answer both problems in the affirmative.

Theorem I (Theorem 5.4, Theorem 7.2)**.** *Let $\mathscr{K}$ be the complement complex constructed from any combinatorial stratification of any essential c-arrangement $\mathscr{A}$ in S^d or $\mathbb{R}^d$. Then $\mathscr{K}$ admits a perfect discrete Morse function.*

Corollary II. *The complement of any affine 2-arrangement in $\mathbb{R}^d$, and the complement of any 2-arrangement in S^d, is a minimal space.*

A crucial step on the way to the proof of Theorem I is the proof of a Lefschetz-type hyperplane theorem for the complements of c-arrangements. The lemma we actually need is a bit technical (Lemma 4.1), but roughly speaking, the result can be phrased in the following way:

Theorem III (Theorem 4.2)**.** *Let $\mathscr{A}^{\mathrm{c}}$ denote the complement of any affine 2-arrangement in $\mathbb{R}^d$, and let H denote a general position hyperplane in $\mathbb{R}^d$. Then $\mathscr{A}^{\mathrm{c}}$ is homotopy equivalent to $H \cap \mathscr{A}^{\mathrm{c}}$ with finitely many e-cells attached, where $e = \lceil d/2 \rceil = d - \lfloor d/2 \rfloor$.*

An analogous theorem holds for complements of c-arrangements (with $e = d - \lfloor d/c \rfloor$); it is an immediate consequence of the analog of Corollary II for c-arrangements, $c \neq 2$, cf. Section 7. Theorem III extends a well known result on complex hyperplane arrangements, which follows from Morse theory applied to the Milnor fiber [Ran02, HL71]. The main ingredients for our proof of Theorems I and III are:

- the formula to compute the homology of subspace arrangements in terms of the intersection lattice, due to Goresky and MacPherson [GM88]; compare Lemma 5.1;
- the study of combinatorial stratifications as initiated by Björner and Ziegler [BZ92]; cf. Section 2.1;
- the study of the collapsibility of complexes endowed with a specific metric, initiated in the papers [AB11] and [AB12b]; this is used in the proof of Theorem 3.1;
- the idea of Alexander duality for Morse functions, in particular the elementary notion of "out-j collapse", introduced in Section 2.2;
- the notion of (Poincaré) duality of discrete Morse functions, which goes back to Forman [For98] and [Ben12]. This is used to establish discrete Morse functions on complement complexes, cf. Theorem 2.3.

2 Preliminaries

We use this section to recall the basic facts on 2-arrangements, combinatorial stratifications and discrete Morse theory. For more on subspace arrangements, we refer the reader to [BZ92, GM88, OT92]. For the dual block complex, we refer to Munkres' book [Mun84, p. 377]. For discrete Morse theory, and in particular dual discrete Morse functions, we refer the reader to Forman [For98] and Benedetti [Ben12]. If H is a hyperplane, and C is a CW-complex embedded in the same space, then H is in *general position* with respect to C if H intersects all elements of C transversally. Similarly, if $\mathscr{A}$ is a subspace arrangement, then H and $\mathscr{A}$ they are in *general position* with respect to each other if H intersects all elements of $\mathscr{A}$, and all intersections of its elements, transversally. To prove the main results, we work with arrangements in S^d, and treat the euclidean case as a special case of the spherical one.

2.1 2-arrangements, combinatorial stratifications, complement complexes

Throughout this paper, d and n will always be non-negative integers. By $[n]$ we denote the set of positive integers that are smaller than or equal to n.

Let S^d be the unit sphere in $\mathbb{R}^{d+1}$. An *i-dimensional subspace in S^d* is the intersection of the unit sphere S^d with some $(i+1)$-dimensional linear subspace of $\mathbb{R}^{d+1}$. A *hyperplane in S^d* is a $(d-1)$-dimensional subspace in S^d. A *c-arrangement $\mathscr{A}$* in S^d (resp. in $\mathbb{R}^d$) is a collection $(h_i)_{i\in[n]}$ of distinct $(d-c)$-dimensional subspaces of S^d (resp. of $\mathbb{R}^d$), such that the codimension of any non-empty intersection of its elements is a multiple of c. In particular, any c-arrangement with $d<c$ is the empty arrangement. A c-arrangement $\mathscr{A}$ is *essential* if the intersection of all the elements of $\mathscr{A}$ is empty, and *non-essential* otherwise. Non-essential arrangements in particular include empty arrangements.

Definition 2.1 (Extensions and stratifications)**.** A *sign extension $\mathscr{A}_\sigma$* of a 2-arrangement $\mathscr{A}=(h_i)_{i\in[n]}$ in S^d is any collection of hyperplanes $(H_i)\subset S^d$, $i\in[n]$ such that for each i, we have $h_i\subset H_i$. We say that the subspaces S^d, H_i and h_i itself *extend* h_i. A *hyperplane extension $\mathscr{A}_e$* of $\mathscr{A}$ in S^d is a sign extension of $\mathscr{A}$ together with an arbitrary collection of hyperplanes in S^d. The elements of $\mathscr{A}_e$ induce a subdivision of S^d into convex subsets; this subdivision $\mathbf{s}(\mathscr{A}_e)$ is the *stratification of S^d given by $\mathscr{A}_e$.*

Definition 2.2 (Fine extensions and combinatorial stratifications)**.** If $\mathscr{A}$ is essential, then any stratification $\mathbf{s}(\mathscr{A}_e)$ of S^d is a regular CW complex. If $\mathscr{A}$ is non-essential, then the stratification $\mathbf{s}(\mathscr{A}_\sigma)$ associated to a sign extension need not be a regular CW complex; however, after adding some extra hyperplanes, some extension $\mathscr{A}_e\supset\mathscr{A}_\sigma$ will give rise to a regular CW complex $\mathbf{s}(\mathscr{A}_e)$. If the extension $\mathscr{A}_e$ gives rise to a stratification $\mathbf{s}(\mathscr{A}_e)$ that *is* a regular CW complex, then $\mathscr{A}_e$ is a *fine* extension and the stratification $\mathbf{s}(\mathscr{A}_e)$ is *combinatorial.*

Let C be a polytopal complex in $\mathbb{R}^d$ (resp. S^d), and let M be an arbitrary subset of $\mathbb{R}^d$ (resp. S^d). The *restriction* $\mathrm{R}(C,M)$ *of C to M* is the maximal subcomplex of C all whose faces are contained in M. Dually, let $\mathbf{s}^*$ be the dual block complex of a combinatorial stratification $\mathbf{s}$ of S^d, and let C be any subcomplex of $\mathbf{s}^*$. Then, for M open, $\mathrm{R}^*(C,M)$ is the subcomplex of $C\subset\mathbf{s}^*$ corresponding to faces of $\mathbf{s}$ intersecting M.

Definition 2.3 (Complement complexes for spherical 2-arrangements)**.** Let $\mathbf{s}$ be any combinatorial stratification of the sphere S^d associated to a 2-arrangement $\mathscr{A}$. Let $\mathbf{s}^*$ be the dual block complex of $\mathbf{s}$. Define $\mathscr{K}(\mathscr{A}_e):=\mathrm{R}^*(\mathbf{s}^*,S^d\setminus\mathscr{A})$. Equivalently, $\mathscr{K}(\mathscr{A}_e)$ is the subcomplex of $\mathbf{s}^*$ consisting of the dual of the faces that are *not* contained in $\mathrm{R}(\mathbf{s}(\mathscr{A}_e),\mathscr{A})$. The CW complex $\mathscr{K}(\mathscr{A}_e)$ is the *complement complex of $\mathscr{A}$ induced by $\mathscr{A}_e$.*

Definition 2.4 (Complement complexes for affine 2-arrangements)**.** Let $\mathscr{A}$ denote a 2-arrangement of *affine* subspaces in $\mathbb{R}^d$, and let ρ denote the radial projection of $\mathbb{R}^d$ to an open hemisphere $\mathcal{D}$ in S^d. The image $\rho(\mathscr{A}):=\{\rho(\sigma):\sigma\in\mathscr{A}\}$ is the restriction of the 2-arrangement $\mathscr{A}':=\{\mathrm{sp}(\sigma)\cap S^d:\sigma\in\rho(\mathscr{A})\}$ in $S^d\subset\mathbb{R}^{d+1}$ to $\mathcal{D}$, where sp denotes the linear span in Euclidean space. Consider the combinatorial stratification $\mathbf{s}(\mathscr{A}'_e)$ of $\mathscr{A}'$ given by a fine extension $\mathscr{A}'_e$, and the complement complex $\mathscr{K}(\mathscr{A}'_e)$ it gives rise to. We say that $\mathscr{K}(\mathscr{A}'_e)$ is the *complement complex of the arrangement $\mathscr{A}$.*

Recall that a *model* for a topological space is a CW complex homotopy equivalent to it.

Lemma 2.5 ([BZ92, Prp. 3.1])**.** *The complement complex of any 2-arrangement $\mathscr{A}$ in $\mathbb{R}^d$ (resp. S^d) is a model for the complement $\mathscr{A}^{\mathrm{c}}=\mathbb{R}^d\setminus\mathscr{A}$ (resp. $S^d\setminus\mathscr{A}$).*

2.2 Outwardly matched faces of subcomplexes; out-j collapses

Definition 2.6 (Outwardly matched faces)**.** Let C be a regular CW complex. Let D be a non-empty subcomplex. Consider a discrete Morse matching $\varPhi$ on C. A face τ of D is *outwardly matched (with respect to the pair* (C, D)*)* if it is matched to a face that does not belong to D.

Definition 2.7 (Out-j collapse of a pair)**.** Let C be a regular CW complex. Let D be a subcomplex. Suppose that C collapses onto a subcomplex C'. The pair $(C,\, D)$ *out-j collapses* to the pair $(C', D \cap C')$, and we write in symbols

$$(C,\, D) \searrow_{\text{out-}j} (C',\, D \cap C')$$

if the collapsing sequence that reduces C to C' can be chosen so that every outwardly matched face of D has dimension j.

Definition 2.8 (Out-j collapsible)**.** Let C be a regular CW complex. Let D be any non-empty subcomplex of C. We say that the pair (C, D) is *out-j collapsible*, if there is a vertex v of D such that

$$(C,\, D) \searrow_{\text{out-}j} (v,\, v).$$

Furthermore, the pair $(C, \emptyset)$ is *out-j collapsible* for every integer j if C is collapsible.

Example 2.9. Let C be a collapsible simplicial d-complex ($d \geq 1$).

- The pairs (C, C) and $(C, \emptyset)$ are out-j collapsible for any j.
- There is a tree D in C such that (C, D) is out-j collapsible for any j. If C triangulates a disk, any tree D has this property.
- If D is the k-skeleton of C, with $0 \leq k < d$, then (C, D) is out-k collapsible.
- If D is a triangulation of the Dunce hat that is a subcomplex of C, then (C, D) is not out-j collapsible for any j. In fact, any collapsing sequence for C pairs some triangle of D with some tetrahedron, and also some edge of D with a triangle outside D, cf. Proposition 2.10.

Proposition 2.10. *Let (C, D) be an out-j collapsible pair, with D non-empty. The number of outwardly matched j-faces of D is independent of the collapsing sequence chosen, and equal to* $(-1)^j \cdot (\chi(D) - 1)$*, where χ denotes the Euler characteristic. Moreover, the following are equivalent:*
(1) *D is contractible;*
(2) *D has no outmatched faces;*
(3) *D is collapsible.*

Definition 2.11. With the above notation, if any of the conditions (i), (ii), or (iii) is satisfied, we say that (C, D) *is a collapsible pair*. The pair $(C, \emptyset)$ is a *collapsible pair* if C is collapsible.

For example, if C is any collapsible complex, (C, C) is a collapsible pair, and there exists a tree D in C such that (C, D) is a collapsible pair.

Proof of Proposition 2.10. Fix an out-j collapsing sequence for (C, D). Let O be the set of outwardly matched faces of D. Let P be the vertex onto which C collapses. Let Q be the set of the faces of D that are matched together with another face of D. Clearly, the sets O, P, Q form a partition of the set of all faces of D. We set

$$f_i := \#\{i\text{-faces of } D\}, \quad o_i := \#\{i\text{-faces in } O\}, \quad q_i := \#\{i\text{-faces in } Q\}.$$

Clearly, $f_0 = o_0 + 1 + q_0$, and $f_i = o_i + q_i$ for $i \geq 1$. By definition of out-j collapsibility, $o_i = 0$ unless $i = j$. In particular, $\sum (-1)^i o_i = (-1)^j o_j$. Now, faces matched together in a collapsing sequence must have consecutive dimension. It follows (by pairwise canceling) that $\sum (-1)^i q_i = 0$. Hence,

$$\chi(D) \;=\; \sum (-1)^i f_i \;=\; 1 + \sum (-1)^i o_i + \sum (-1)^i q_i \;=\; 1 + (-1)^j o_j.$$

Multiplying by $(-1)^j$, we obtain

$$(-1)^j \chi(D) = (-1)^j + (-1)^{2j} o_j = (-1)^j + o_j.$$

Hence $o_j = (-1)^j \cdot (\chi(D) - 1)$ and the first claim is proven. For the second part:

- (1) $\Rightarrow$ (2) : If D is contractible, $\chi(D) = 1$. By the formula above, $o_j = 0$.
- (2) $\Rightarrow$ (3) : By assumption, there is a collapsing sequence for C which removes all faces of D in pairs. Consider the restriction to D of the collapsing sequence for C; this yields a collapsing sequence for D.
- (3) $\Rightarrow$ (1) : This is implied by the fact if $D \searrow D'$, then D' is a deformation retract of D. $\square$

2.3 Complement matchings

Let $\mathbf{s} = \mathbf{s}(\mathscr{A}_e)$ denote a combinatorial stratification of the sphere S^d. To obtain Morse matchings on the complement complex $\mathscr{K}(\mathscr{A}_e)$, we first study matchings on the stratification $\mathbf{s}$. Via duality, Morse matchings on $\mathbf{s}$ will then give rise to Morse matchings on $\mathscr{K}$.

Definition 2.12 (Restrictions of matchings)**.** Let Φ be any discrete Morse matching on a regular CW complex C such that, for some subcomplex D of C, the Morse matching Φ contains no outwardly matched faces with respect to the pair (C, D). Let us denote by Φ_D the restriction of Φ to D, that is, the collection of all matchings in Φ involving two faces of D.

Definition 2.13 (Complement matching)**.** Consider the dual Morse matching Φ^* on $\mathbf{s}^*$ of the Morse matching Φ defined on a combinatorial stratification $\mathbf{s}$. The matching Φ^* has an outwardly matched face σ^* (matched with a face τ^*) with respect to the pair $(\mathbf{s}^*, \mathscr{K})$ for every outwardly matched face τ (matched with σ) of Φ with respect to the pair $(\mathbf{s}, \mathrm{R}(\mathbf{s}, \mathscr{A}))$. After we remove all such matchings from Φ^*, we are left with a Morse matching on $\mathbf{s}^*$ that has no outwardly matched faces with respect to the pair $(\mathbf{s}^*, \mathscr{K})$. After we restrict this matching to the faces of $\mathscr{K}$, we obtain a matching on the complex $\mathscr{K}$, the *complement matching* Φ^{c} of Φ.



Figure 1: An outward matching (σ^*, τ^*) of the pair $(\mathbf{s}^*, \mathscr{K})$ corresponds to an outward matching (τ, σ) of the pair $(\mathbf{s}, \mathrm{R}(\mathbf{s}, \mathscr{A}))$.

These notions allow us to study Morse matchings on the complement complex by studying Morse matchings on the stratification itself.

Theorem 2.14. *Let $\mathscr{A}$ be an arbitrary 2-arrangement. Let $\mathbf{s} = \mathbf{s}(\mathscr{A}_e)$ be any combinatorial stratification of S^d. Consider a Morse matching Φ on $\mathbf{s}$. Then the critical i-faces of Φ^{c} are in one-to-one correspondence with the union of*
- *the critical $(d-i)$-faces of Φ that are not faces of $\mathrm{R}(\mathbf{s}, \mathscr{A})$, and*
- *the outwardly matched $(d-i-1)$-faces of Φ with respect to the pair $(\mathbf{s}, \mathrm{R}(\mathbf{s}, \mathscr{A}))$.*

If M is a subset of S^d such that all noncritical faces of Φ intersect M, then the critical i-faces of $\Phi^{\mathrm{c}}_{\mathrm{R}^*(\mathscr{K},M)}$ are in bijection with the union of
- *the critical $(d-i)$-faces of Φ that are not faces of $\mathrm{R}(\mathbf{s}, \mathscr{A})$, and that intersect M, and*
- *the outwardly matched $(d-i-1)$-faces of Φ with respect to the pair $(\mathbf{s}, \mathrm{R}(\mathbf{s}, \mathscr{A}))$.*

Proof. Both bijections are natural: Each critical i-face of Φ^{c} corresponds to either a critical i-face of Φ^* or to an outwardly-matched i-face of Φ^* with respect to the pair $(\mathbf{s}^*, \mathscr{K})$. Now,
- critical i-faces of Φ^* in $\mathscr{K}$ are in one-to-one correspondence with the critical $(d-i)$-faces of Φ that are not faces of $\mathrm{R}(\mathbf{s}, \mathscr{A})$, and
- the outwardly matched i-faces of Φ^* with respect to the pair $(\mathbf{s}^*, \mathscr{K})$, are in one-to-one correspondence with the outwardly matched $(d-i-1)$-faces of Φ with respect to the pair $(\mathbf{s}, \mathrm{R}(\mathbf{s}, \mathscr{A}))$.

Since the two sets are clearly disjoint, we obtained the desired bijection for the critical faces of Φ^{c}. The bijection for the critical faces of $\Phi^{\mathrm{c}}_{\mathrm{R}^*(\mathscr{K},M)}$ is obtained analogously. □

3 Restricting stratifications to general position hemispheres

In this section, we study Morse matchings on a stratification of a 2-arrangement in S^d. More precisely, we study Morse matchings on the restrictions of a stratification to an open hemisphere. The main result of this section is Theorem 3.1, which will turn out to be crucial in order to establish Main Theorem I.

We say that a hemisphere is in *general position* with respect to an object (a CW-complex, a subspace arrangement) if the boundary of the hemisphere is in general position with respect to the object. If $\mathscr{A}$ is any subspace arrangement in S^d, and H is a subspace of S^d, we define

$$\mathscr{A}^H := \{\sigma \cap H : \sigma \in \mathscr{A}\}.$$

Let $T_p^1 S^d$ denote the unit ball in the tangent space $T_p S^d$ of S^d at a point p in S^d. If X is any convex subset of S^d, define $T_p^1 X := T_p^1 S^d \cap T_p X$. If $\mathscr{A}$ is any subspace arrangement in S^d, we define the *link* $\mathrm{Lk}(p, \mathscr{A})$ of $\mathscr{A}$ at p by

$$\mathrm{Lk}(p, \mathscr{A}) := \{T_p^1 \sigma : \sigma \in \mathscr{A},\ \sigma \cap p \neq \emptyset\} \subset T_p^1 S^d.$$

Similarly, if C is a regular CW complex in S^d all whose cells are convex (e.g. a subcomplex of a stratification of S^d), and v is a vertex, the *link* $\mathrm{Lk}(v, C)$ of C at v is the regular CW complex

$$\mathrm{Lk}(v, C) := \{T_v^1 \sigma : \sigma \in C,\ v \in \sigma\} \subset T_v^1 S^d.$$

Our first goal is to investigate whether, for any given hemisphere $\mathcal{D}$, the pair $(\mathrm{R}(\mathbf{s}, \mathcal{D}), \mathrm{R}(\mathbf{s}, \mathcal{D} \cap \mathscr{A}))$ is out-j collapsible for some suitable integer j. As it turns out, the right j to consider is the integer

$$\iota(d) := \lfloor d/2 \rfloor - 1.$$

Theorem 3.1. *Let $\mathscr{A}$ be any 2-arrangement. Let $\mathbf{s} = \mathbf{s}(\mathscr{A}_e)$ be a combinatorial stratification of S^d. Let $\mathcal{D}$ be a closed hemisphere in general position with respect to $\mathbf{s}(\mathscr{A}_e)$. We have:*

(A) *For any nonempty k-dimensional subspace H of $\mathscr{A}_\sigma \subset \mathscr{A}_e$ extending an element of $\mathscr{A}$, the pair $(\mathrm{R}(\mathbf{s}, \mathcal{D} \cap H), \mathrm{R}(\mathbf{s}, \mathcal{D} \cap \mathscr{A} \cap H))$ is out-$\iota(d)$ collapsible.*

(B) *If $\mathscr{A}$ is non-essential, then $(\mathrm{R}(\mathbf{s}, \mathcal{D}), \mathrm{R}(\mathbf{s}, \mathcal{D} \cap \mathscr{A}))$ is a collapsible pair.*

Proof. To simplify the notation, for any subset M of S^d we set

$$\mathrm{R}\,[M] := \mathrm{R}\,(\mathbf{s}(\mathscr{A}_e),\, M) \quad \text{and} \quad \mathrm{R}'\,[M] := \mathrm{R}\,(\mathbf{s}(\mathscr{A}_e),\, M \cap \mathscr{A}) \subset \mathrm{R}\,[M].$$

We proceed by induction on d and k. Let $A_{d,k}$ denote the statement that part (A) is proven for all spheres of dimension d and subspaces H of dimension k. Here, k satisfies $d \geq k \geq \max\{d-2,\, 0\}$, since H is nonempty and extends an element of $\mathscr{A}$. Similarly, let B_d denote the statement that part (B) holds for arrangements in spheres of dimension d.

For the base cases of the induction, it suffices to treat the case $A_{0,0}$. In this case $\mathrm{R}\,[\mathcal{D}] = \mathrm{R}\,[\mathcal{D} \cap H]$ is a vertex. Thus, there is no further collapse of $\mathrm{R}\,[\mathcal{D} \cap H]$ necessary, proving that $(\mathrm{R}\,[\mathcal{D} \cap H], \mathrm{R}'\,[\mathcal{D} \cap H])$ is a collapsible pair, in particular out-$\iota(d)$ collapsible.

Our inductive proof proceeds like this:

I. We prove that $A_{d,d}$ implies B_d for $d \geq 0$.

II. We prove that $A_{k,k}$ implies $A_{d,k}$ for $k = \max\{d-2,\, 0\}$.

III. We prove that B_{k-1}, $A_{k-1,k-1}$ and $A_{d,k-1}$ together imply $A_{d,k}$ for $k > \max\{d-2,\, 0\}$.

Part I. $A_{d,d}$ **implies** B_d **for** $d \geq 0$

By the second part of Proposition 2.10 and assumption $A_{d,d}$, we only have to prove that $\mathrm{R}'[\mathcal{D}]$ is contractible or empty. Let σ denote the intersection of all elements of $\mathscr{A}$. If $\mathscr{A}$ is nonempty, then so is σ and $\mathrm{R}'[\mathcal{D}]$ deformation retracts onto the contractible complex $\mathrm{R}[\mathcal{D} \cap \sigma]$. If $\mathscr{A}$ is empty, then so is $\mathrm{R}'[\mathcal{D}]$.

Part II. $A_{k,k}$ **implies** $A_{d,k}$ **for** $k = \max\{d-2,\, 0\}$

We have to show that $(\mathrm{R}\,[\mathcal{D} \cap H],\, \mathrm{R}'\,[\mathcal{D} \cap H])$ is out-$\iota(d)$ collapsible. We will show something stronger, namely, that it is a collapsible pair. If $k = 0$, then $\mathrm{R}\,[\mathcal{D} \cap H]$ is a vertex, thus $(\mathrm{R}\,[\mathcal{D} \cap H],\, \mathrm{R}'\,[\mathcal{D} \cap H])$ is a collapsible pair by definition.

Thus, we may assume $k = d - 2$. The k-dimensional subspace H is an element of $\mathscr{A}$, so $\mathrm{R}\,[\mathcal{D} \cap H]$ lies in $\mathscr{A}$, we have

$$(\mathrm{R}\,[\mathcal{D} \cap H],\, \mathrm{R}'\,[\mathcal{D} \cap H]) = \big(\mathrm{R}(\mathbf{s},\, \mathcal{D} \cap H)\,,\, \mathrm{R}(\mathbf{s},\, \mathcal{D} \cap H \cap \mathscr{A})\big) = \big(\mathrm{R}(\mathbf{s},\, \mathcal{D} \cap H)\,,\, \mathrm{R}(\mathbf{s},\, \mathcal{D} \cap H)\big).$$

But a pair (C, C), C a regular CW complex, is a collapsible pair iff C is collapsible. By definition, if a pair is out-$\iota(d)$ collapsible, the first complex in the pair is collapsible; so, since the pair $(\mathrm{R}[\mathcal{D}\cap H], \mathrm{R}[\mathcal{D}\cap H\cap(\mathscr{A}\setminus H)])$ is out-$\iota(d)$ collapsible by assumption $A_{k,k}$, it follows trivially that $\mathrm{R}[\mathcal{D}\cap H]$ is collapsible.

Part III. B_{k-1}**,** $A_{k-1,k-1}$ **and** $A_{d,k-1}$ **together imply** $A_{d,k}$ **for** $k-1\geq\max\{d-2,\,0\}$

Let h denote an element of $\mathscr{A}$ extended by H. Let $\eta\in\mathscr{A}_e$ be a codimension-one subspace of H that extends h as well. Let η^+ be a closed hemisphere in H delimited by η, and let η^- denote the closed hemisphere complementary to η^+. We need to prove that the pair $(\mathrm{R}\,[\mathcal{D}\cap H],\,\mathrm{R}'\,[\mathcal{D}\cap H])$ is out-$\iota(d)$ collapsible. The proof consists of three steps:

(1) We prove $(\mathrm{R}\,[\mathcal{D}\cap H],\,\mathrm{R}'\,[\mathcal{D}\cap H])\searrow_{\text{out-}\iota(d)}(\mathrm{R}\,[\mathcal{D}\cap\eta^-],\,\mathrm{R}'\,[\mathcal{D}\cap\eta^-])$.

(2) Symmetrically, we prove $(\mathrm{R}\,[\mathcal{D}\cap\eta^-],\,\mathrm{R}'\,[\mathcal{D}\cap\eta^-])\searrow_{\text{out-}\iota(d)}(\mathrm{R}\,[\mathcal{D}\cap\eta],\,\mathrm{R}'\,[\mathcal{D}\cap\eta])$.

(3) It then remains to show that $(\mathrm{R}\,[\mathcal{D}\cap\eta],\,\mathrm{R}'\,[\mathcal{D}\cap\eta])$ is out-$\iota(d)$ collapsible. This, however, is true by assumption $A_{d,k-1}$.

Step (2) is completely analogous step (1), so it is left out. It remains to prove step (1); we show

$$(\mathrm{R}\,[\mathcal{D}\cap H],\,\mathrm{R}'\,[\mathcal{D}\cap H])\searrow_{\text{out-}\iota(d)}(\mathrm{R}\,[\mathcal{D}\cap\eta^-],\,\mathrm{R}'\,[\mathcal{D}\cap\eta^-]).$$

To achieve this, we establish a geometry-based total order on the vertices of $\mathrm{R}\,[\mathcal{D}\cap H]$, and we collapse them away one at the time. In details: Let π denote a central projection of $\mathcal{D}\cap H$ to $\mathbb{R}^{d-1}$, and let $\vec{n}^+$ denote the inner normal to $\pi(\eta^+)\subset\mathbb{R}^{d-1}$. Perturb $\vec{n}^+$ to a vector $\vec{n}$ such that, for any two vertices v, w of $\mathrm{R}\,[\mathcal{D}\cap H]$,

- if $\langle\,\pi(v),\vec{n}^+\,\rangle\;>\;\langle\,\pi(w),\,\vec{n}^+\,\rangle$, then $\langle\,\pi(v),\,\vec{n}\,\rangle\;>\;\langle\,\pi(w),\,\vec{n}\,\rangle$;
- if $\langle\,\pi(v),\vec{n}\,\rangle\;=\;\langle\,\pi(w),\,\vec{n}\,\rangle$, then $v\;=\;w$.

The function

$$x\;\longmapsto\;\langle\,\pi(x),\,\vec{n}\,\rangle$$

induces a total order on the vertices of $\mathrm{R}\,[\mathcal{D}\cap H]$, starting with the vertex with the highest value under this function.

Let v be the vertex of $\mathrm{R}\,[\mathcal{D}\cap H]$ maximizing the function $x\;\mapsto\;\langle\,\pi(x),\,\vec{n}\,\rangle$. If v is a vertex of $\mathrm{R}\,[\mathcal{D}\cap\eta^-]$, then $\mathrm{R}\,[\mathcal{D}\cap H]=\mathrm{R}\,[\mathcal{D}\cap\eta^-]$ and we are done. If not, notice that $\mathrm{Lk}(v,\mathbf{s}(\mathscr{A}_e^H))$ is a combinatorial stratification of the 2-arrangement $\mathrm{Lk}(v,\mathscr{A}^H)$ in the $(k-1)$-sphere T_v^1H, induced by the hyperplane extension $\mathrm{Lk}(v,\mathscr{A}_e^H)$ of $\mathrm{Lk}(v,\mathscr{A}^H)$. The complex $\mathrm{Lk}(v,\mathrm{R}\,[\mathcal{D}\cap H])$ is the restriction of $\mathrm{Lk}(v,\mathbf{s}(\mathscr{A}_e^H))$ to the general position hemisphere $T_v^1\mathcal{D}_v$, where

$$\mathcal{D}_v\;:=\;\pi^{-1}\{\,x\in\mathbb{R}^{k-1}:\langle\,\pi(v),\,\vec{n}\,\rangle\;>\;\langle\,x,\,\vec{n}\,\rangle\},$$

since v maximizes $\langle\,\pi(x),\,\vec{n}\,\rangle$. At this point, we want to apply the induction assumptions $A_{k-1,k-1}$ and B_{k-1} to $\mathrm{Lk}(v,\mathrm{R}\,[\mathcal{D}\cap H])$. There are two cases to consider: either k is divisible by 2, or not.

- If $k\;=\;1$ mod 2, then $\mathrm{Lk}(v,\mathscr{A}^H)$ is a *non-essential* 2*-arrangement* in T_v^1H. Thus, by assumption (B_{d-1}), the pair

  $$\big(\mathrm{Lk}(v,\mathrm{R}[\mathcal{D}\cap H]),\mathrm{Lk}(v,\mathrm{R}'[\mathcal{D}\cap H])\big),$$

  is a collapsible pair. Consequently, $(\mathrm{R}\,[\mathcal{D}\cap H],\mathrm{R}'\,[\mathcal{D}\cap H])$ collapses to $(\mathrm{R}\,[\mathcal{D}\cap H]-v,\mathrm{R}'\,[\mathcal{D}\cap H]-v)$ without any new outwardly matched faces (here, we use the minus sign to denote the deletion of a subcomplex). In particular,

  $$(\mathrm{R}\,[\mathcal{D}\cap H],\mathrm{R}'\,[\mathcal{D}\cap H])\searrow_{\text{out-}\iota(d)}(\mathrm{R}\,[\mathcal{D}\cap H]-v,\mathrm{R}'\,[\mathcal{D}\cap H]-v).$$

- If $k=0$ mod 2, then $\mathrm{Lk}(v,\mathscr{A}^H)$ is a 2-arrangement in T_v^1H. Thus, by assumption $(A_{k-1,k-1})$, the pair

  $$\big(\mathrm{Lk}(v,\mathrm{R}[\mathcal{D}\cap H]),\mathrm{Lk}(v,\mathrm{R}'[\mathcal{D}\cap H])\big),$$

  is an out-$\iota(k-1)$ collapsible pair. We have $\iota(k-1)=\iota(k)-1$ (since $k=0$ mod 2) and $\iota(k)=\iota(d)$ (since $d-2\leq k\leq d$), so that $\iota(k-1)=\iota(d)-1$. Hence we obtain that the pair $(\mathrm{R}\,[\mathcal{D}\cap H],\mathrm{R}'\,[\mathcal{D}\cap H])$ out-$\iota(d)$ collapses to $(\mathrm{R}\,[\mathcal{D}\cap H]-v,\mathrm{R}'\,[\mathcal{D}\cap H]-v)$.

If there are no more vertices of $\mathrm{R}\,[\mathcal{D}\cap H]-v$ in the interior of η^+, we are done with the proof of step (1). If not, we repeat the above procedure, this time using $(\mathrm{R}\,[\mathcal{D}\cap H]-v,\mathrm{R}'\,[\mathcal{D}\cap H]-v)$ instead of $(\mathrm{R}\,[\mathcal{D}\cap H],\mathrm{R}'\,[\mathcal{D}\cap H])$, and the vertex w of $\mathrm{R}\,[\mathcal{D}\cap H]-v$ that maximizes $\langle\,\pi(x),\,\vec{n}\,\rangle$ instead of v. This is continued until all vertices in $\mathrm{R}(\mathbf{s}(\mathscr{A}_e),\mathcal{D}\cap\mathrm{int}(\eta^+))$ are deleted, which shows that the pair $(\mathrm{R}\,[\mathcal{D}\cap H],\,\mathrm{R}'\,[\mathcal{D}\cap H])$ can be out-$\iota(d)$ collapsed to $(\mathrm{R}\,[\mathcal{D}\cap\eta^-],\,\mathrm{R}'\,[\mathcal{D}\cap\eta^-])$. $\square$

4 Proof of Theorem III

We have now all the tools to prove our Lefschetz-type hyperplane theorem (Theorem 4.2). First, a lemma:

Lemma 4.1. *Let* $\mathbf{s}(\mathscr{A}_e)$ *denote any combinatorial stratification of* S^d *associated to any* 2*-arrangement* $\mathscr{A}$*, and let* $\mathcal{D}$*,* $\mathcal{D}'$ *denote two closed hemispheres such that* $\mathcal{D}'$ *is in general position with respect to* $\mathbf{s}(\mathscr{A}_e \cup \partial\mathcal{D})$*. Then*

$$\mathrm{R}(\mathbf{s}, \mathcal{D} \cup \mathcal{D}') \searrow_{\textit{out-}\iota(d)} \mathrm{R}(\mathbf{s}, \mathcal{D}).$$

Proof. We may assume that $\mathcal{D} \setminus \mathcal{D}'$ contains a single vertex of $\mathbf{s}(\mathscr{A}_e)$; if this is not immediately the case, perturb $\mathcal{D}'$ slightly and find a sequence of hemispheres $\mathcal{D}_0 = \mathcal{D}, \mathcal{D}_1, \ldots, \mathcal{D}_n = \mathcal{D}'$ such that

- $\mathcal{D} \cap \mathcal{D}' \subset \mathcal{D}_i$ for all i from 0 to n,
- $\mathcal{D}_{i+1} \setminus \mathcal{D}_i$ contains only one vertex of $\mathbf{s}(\mathscr{A}_e)$, for each $i \in \{0, \ldots, n-1\}$, and
- $\mathcal{D}_{i+1}$, $i \in [n-1]$ is in general position with respect to $\mathbf{s}(\mathscr{A}_e \cup \partial\mathcal{D}_i)$.

A simple iteration of the case where a single vertex of $\mathbf{s} = \mathbf{s}(\mathscr{A}_e)$ is in $\mathcal{D} \setminus \mathcal{D}'$ then gives the desired result. Thus, let v denote the vertex of $\mathbf{s}$ in $\mathcal{D} \setminus \mathcal{D}'$. Let $\mathcal{D}_v$ denote a hemisphere in S^d such that $\mathcal{D} \cap \mathcal{D}' \subset \mathcal{D}_v$ and such that v lies in the boundary of $\mathcal{D}_v$. Then $\mathrm{Lk}(v, \mathbf{s})$ is a combinatorial stratification of the 2-arrangement $\mathrm{Lk}(v, \mathscr{A})$ in the $(d-1)$-sphere $T^1_v S^d$ and the complex $\mathrm{Lk}(v, \mathrm{R}(\mathbf{s}, \mathcal{D} \cup \mathcal{D}'))$ is the restriction of $\mathrm{Lk}(v, \mathbf{s})$ to the general position hemisphere $T^1_v \mathcal{D}_v$. Thus, as in Part III of Theorem 3.1, there are two cases:

- If $d \neq 0 \bmod 2$, then $\mathrm{Lk}(v, \mathscr{A})$ is a *non-essential* 2-arrangement in $T^1_v S^d$. Thus, by Theorem 3.1(B), the pair
  $$\big(\mathrm{Lk}(v, \mathrm{R}(\mathbf{s}, \mathcal{D})), \mathrm{Lk}(v, \mathrm{R}(\mathbf{s}, \mathcal{D} \cap \mathscr{A}))\big),$$
  is a collapsible pair. Consequently, the pair $\big(\mathrm{R}(\mathbf{s}, \mathcal{D} \cup \mathcal{D}'), \mathrm{R}(\mathbf{s}, (\mathcal{D} \cup \mathcal{D}') \cap \mathscr{A})\big)$ collapses to the pair
  $$\big(\mathrm{R}(\mathbf{s}, \mathcal{D} \cup \mathcal{D}') - v, \mathrm{R}(\mathbf{s}, (\mathcal{D} \cup \mathcal{D}') \cap \mathscr{A}) - v\big) = \big(\mathrm{R}(\mathbf{s}, \mathcal{D}), \mathrm{R}(\mathbf{s}, \mathcal{D} \cap \mathscr{A})\big)$$
  without any new outwardly matched faces.
- If $d = 0 \bmod 2$, then $\mathrm{Lk}(v, \mathscr{A})$ is a 2-arrangement in the $T^1_v S^d$. By Theorem 3.1(A), the pair
  $$\big(\mathrm{Lk}(v, \mathrm{R}(\mathbf{s}, \mathcal{D})), \mathrm{Lk}(v, \mathrm{R}(\mathbf{s}, \mathcal{D} \cap \mathscr{A}))\big),$$
  is an out-$\iota(d-1)$ collapsible pair. Since we have $\iota(d-1) = \iota(d) - 1$, we obtain that the pair $\big(\mathrm{R}\,(\mathbf{s}, \mathcal{D} \cup \mathcal{D}'), \mathrm{R}'\,(\mathbf{s}, (\mathcal{D} \cup \mathcal{D}') \cap \mathscr{A})\big)$ out-$\iota(d)$ collapses to $\big(\mathrm{R}(\mathbf{s}, \mathcal{D}), \mathrm{R}(\mathbf{s}, \mathcal{D} \cap \mathscr{A})\big)$. ☐

Theorem 4.2. *Consider any affine* 2*-arrangement* $\mathscr{A}$ *in* $\mathbb{R}^d$*, and let* H *denote a hyperplane in* $\mathbb{R}^d$ *in general position with respect to* $\mathscr{A}$*. Then the complement* $\mathscr{A}^{\mathrm{c}}$ *of* $\mathscr{A}$ *is homotopy equivalent to* $H \cap \mathscr{A}^{\mathrm{c}}$ *with* e*-cells attached to it, where* $e = \lceil d/2 \rceil = d - \lfloor d/2 \rfloor$*.*

Proof. Define ρ, $\mathcal{D}$ and $\mathscr{A}'$ in S^d in Definition 2.4, and define the hyperplane $H_\rho = \mathrm{sp}(\rho(H)) \cap S^d$ induced by H in $\mathbb{R}^d$. Let $\mathcal{D}^{\mathrm{c}}$ denote the closed hemisphere complementary to $\mathcal{D}$. Let $\mathbf{s}$ be any combinatorial stratification of S^d induced by an hyperplane extension of $\mathscr{A}'$. By Lemma 4.1, $\mathbf{s}$ out-$\iota(d)$ collapses to $\mathrm{R}(\mathbf{s}, \mathcal{D}^{\mathrm{c}})$ after the faces intersecting $H_\rho \cap \mathcal{D}$ are removed. Equivalently, there exists a discrete Morse function $\varPhi$ on $\mathbf{s}$ with the following properties:

- the critical faces of $\varPhi$ are the faces of $\mathrm{R}(\mathbf{s}, \mathcal{D}^{\mathrm{c}})$ and the faces of $\mathbf{s}$ that intersect $H_\rho \cap \mathcal{D}$;
- the outwardly-matched faces of the pair $(\mathbf{s}, \mathrm{R}(\mathbf{s}, \mathscr{A}))$ are all of dimension $\iota(d)$.

By Theorem 2.14 the complementary matching $\varPhi^{\mathrm{c}}_{\mathrm{R}^*(\mathscr{K}(\mathscr{A}'), \mathcal{D})}$ restricted to the faces of $\mathrm{R}^*(\mathscr{K}(\mathscr{A}'), \mathcal{D}) \cong \mathscr{K}(\mathscr{A})$ has the following critical faces:

- the faces of $\mathrm{R}^*(\mathscr{K}(\mathscr{A}'), \mathcal{D} \cap H_\rho)$, whose duals are critical in $(\mathbf{s}, \mathrm{R}(\mathbf{s}, \mathscr{A}))$, and
- the critical faces of dimension $e = \lceil d/2 \rceil = d - \iota(d) - 1$, which correspond to the outwardly matched faces of $\varPhi$.

Thus, $\mathrm{R}^*(\mathscr{K}(\mathscr{A}'), \mathcal{D})$ collapses to $\mathrm{R}^*(\mathscr{K}(\mathscr{A}'), \mathcal{D} \cap H_\rho)$ after the removal of critical cells of dimension e. Since $\mathrm{R}^*(\mathscr{K}(\mathscr{A}'), \mathcal{D}) \simeq \mathscr{K}(\mathscr{A}) \simeq \mathbb{R}^d \setminus \mathscr{A}$ and $\mathrm{R}^*(\mathscr{K}(\mathscr{A}'), \mathcal{D} \cap H_\rho) \simeq H \setminus \mathscr{A}$, we obtain that $\mathbb{R}^d \setminus \mathscr{A}$ is homotopy equivalent to $H \setminus \mathscr{A}$ with e-dimensional cells attached to it. ☐

5 Proof of Theorem I

We start with an easy consequence of the formula of Goresky–MacPherson. For completeness, we provide the (straightforward) proof in the appendix.

Lemma 5.1. *Let $\mathscr{A}$ denote any subspace arrangement in S^d, and let $\mathcal{D}$ denote any open hemisphere of S^d, and let H be any hyperplane in S^d. Then we have*

(I) *If $\mathcal{D}$ is in general position with respect to $\mathscr{A}$, then $\beta_i(S^d \setminus \mathscr{A}) \geq \beta_i(\mathcal{D} \setminus \mathscr{A})$ for all i.*

(II) *If H is in general position with respect to $\mathscr{A}$ and $\mathcal{D}$, then $\beta_i(\mathcal{D} \setminus \mathscr{A}) \geq \beta_i((\mathcal{D} \cap H) \setminus \mathscr{A})$ for all i.*

Besides this lemma, we need the following concept. For any convex set σ and any hyperplane H in S^d, let us denote by σ^H the intersection of σ with H. If C is any subcomplex of a combinatorial stratification of S^d, we also define $C^H := \bigcup_{\sigma \in C} \sigma^H$.

Example 5.2 (Lifting a Morse Matching)**.** Let σ be a convex cell in S^d. This means σ is the intersection of finitely many hemispheres, with the property that the faces of σ form a regular CW complex. Let H denote a general position hyperplane in S^d. Then σ^H is a convex cell in H, and if τ^H is a facet of σ^H, then there exists a unique facet τ of σ with the property that $\tau^H := \tau \cap H$.

Consider now any subcomplex C of a combinatorial stratification of S^d, and any Morse matching φ on the faces of C^H. Then we can match τ with σ for every matched pair of faces τ^H, σ^H in the matching φ of C^H. This gives rise to a Morse matching $\varPhi$ of the faces of C intersecting H from any Morse matching φ on C^H, the *lift* of φ to C.



Figure 2: The lift of a matching.

Lemma 5.3. *Let $\mathbf{s} = \mathbf{s}(\mathscr{A}_e)$ be any stratification of any 2-arrangement $\mathscr{A}$ in S^d induced by any fine extension $\mathscr{A}_e$ of $\mathscr{A}$. Let $\mathcal{D}$ denote any closed hemisphere of S^d, and let $\mathcal{D}^{\mathrm{c}} := S^d \setminus \mathcal{D}$ be its complement. Then, there exists a discrete Morse function X on $\mathbf{s}$ whose critical faces are the subcomplex $\mathrm{R}(\mathbf{s}, \mathcal{D})$ and some additional facet of $\mathbf{s}$ such that the complement matching $X^{\mathrm{c}}_{\mathrm{R}^*(\mathscr{K}, \mathcal{D}^{\mathrm{c}})}$ is perfect on $\mathrm{R}^*(\mathscr{K}, \mathcal{D}^{\mathrm{c}})$.*

Proof. We abbreviate $\mathrm{R}^*[M] := \mathrm{R}^*(\mathscr{K}, M)$ for any set M. The Morse complex of a discrete Morse function is denoted by $\Sigma(\cdot)$. We prove the theorem by induction on the dimension; The case of $d = 0$ is clearly true, since in this case $\mathrm{R}(\mathbf{s}, \mathcal{D})$ is just a vertex. Let H denote any generic hyperplane in S^d. By induction on the dimension, there exists a discrete Morse function φ on $\mathbf{s}^H$ such that the matching $\varphi^{\mathrm{c}}_{\mathrm{R}^*(\mathscr{K}, \mathcal{D}^{\mathrm{c}} \cap H)}$ is perfect on $\mathrm{R}^*(\mathscr{K}, \mathcal{D}^{\mathrm{c}} \cap H)$. We lift φ to a Morse Matching $\varPhi$ of the faces of $\mathbf{s}$ intersecting H. The matching $\varPhi$ collapses, after the removal of a facet F, the combinatorial stratification $\mathbf{s}$ to the subcomplex $\mathrm{R}(\mathbf{s}, S^d \setminus (H \cap \mathcal{D}^{\mathrm{c}}))$.

Now, by Lemma 4.1, $\mathrm{R}(\mathbf{s}, S^d \setminus (H \cap \mathcal{D}^{\mathrm{c}}))$ out-$\iota(d)$ collapses to $\mathrm{R}(\mathbf{s}, \mathcal{D})$. Denote the collapsing sequence demonstrating this collapse by X. We claim that the union $\varPsi$ of the Morse matchings $\varPhi$ and X gives the desired collapse of $\mathbf{s}$ (minus a facet) to $\mathrm{R}(\mathbf{s}, \mathcal{D})$. Clearly, $\varPsi$ induces the collapse

$$\mathbf{s} - F \searrow \mathrm{R}(\mathbf{s}, S^d \setminus (H \cap \mathcal{D}^{\mathrm{c}})) \searrow_{\text{out-}\iota(d)} \mathrm{R}(\mathbf{s}, \mathcal{D}).$$

It remains to demonstrate that $\varPsi^{\mathrm{c}}_{\mathrm{R}^*[\mathcal{D}^{\mathrm{c}}]}$ is perfect. First of all, $\Sigma(\varPhi^{\mathrm{c}}_{\mathrm{R}^*[\mathcal{D}^{\mathrm{c}} \cap H]})$ is homotopy equivalent to $(\mathcal{D}^{\mathrm{c}} \cap H) \setminus \mathscr{A}$, so

$$f_i\big(\Sigma(\varPhi^{\mathrm{c}}_{\mathrm{R}^*[\mathcal{D}^{\mathrm{c}} \cap H]})\big) = c_i(\varPhi^{\mathrm{c}}_{\mathrm{R}^*[\mathcal{D}^{\mathrm{c}} \cap H]}) = \beta_i\big((\mathcal{D}^{\mathrm{c}} \cap H) \setminus \mathscr{A}\big) \text{ for all } i.$$

In particular, $\Sigma(\mathit{\Phi}^{\mathrm{c}}_{\mathrm{R}^*[\mathcal{D}^{\mathrm{c}}\cap H]})$ is a minimal model for $(\mathcal{D}^{\mathrm{c}}\cap H)\setminus\mathscr{A}$. Furthermore, since $\mathrm{R}(\mathbf{s}, S^d\setminus(H\cap\mathcal{D}^{\mathrm{c}}))$ out-$\iota(d)$ collapses to $\mathrm{R}(\mathbf{s},\mathcal{D})$, the Morse complex $\Sigma(\mathit{\Psi}^{\mathrm{c}}_{\mathrm{R}^*[\mathcal{D}^{\mathrm{c}}]})$ is obtained from the complex $\Sigma(\mathit{\Phi}^{\mathrm{c}}_{\mathrm{R}^*[\mathcal{D}^{\mathrm{c}}\cap H]})$ by attaching cells of dimension $e=\lceil d/2\rceil = d-\iota(d)-1$. The attachment of each of these cells contributes to the homology of $\Sigma(\mathit{\Psi}^{\mathrm{c}}_{\mathrm{R}^*[\mathcal{D}^{\mathrm{c}}]})$ either by deleting a generator of the homology in dimension $e-1$, or by adding a generator for the homology in dimension e. But if some of the e-cells deletes a generator in homology, then

$$\beta_{e-1}\big(\mathcal{D}^{\mathrm{c}}\setminus\mathscr{A}\big) = \beta_{e-1}\big(\Sigma(\mathit{\Psi}^{\mathrm{c}}_{\mathrm{R}^*[\mathcal{D}^{\mathrm{c}}]})\big) < \beta_{e-1}\big(\Sigma(\mathit{\Phi}^{\mathrm{c}}_{\mathrm{R}^*[\mathcal{D}^{\mathrm{c}}\cap H]})\big) = \beta_{e-1}\big((\mathcal{D}^{\mathrm{c}}\cap H)\setminus\mathscr{A}\big),$$

in contradiction with Lemma 5.1(II).

Thus, every e-cell attached must add a generator in homology. Since $c_i(\mathit{\Phi}^{\mathrm{c}}_{\mathrm{R}^*[\mathcal{D}^{\mathrm{c}}\cap H]}) = \beta_i\big((\mathcal{D}^{\mathrm{c}}\cap H)\setminus\mathscr{A}\big)$ for all i, we have $c_i(\mathit{\Psi}^{\mathrm{c}}_{\mathrm{R}^*[\mathcal{D}^{\mathrm{c}}]}) = \beta_i\big(\mathcal{D}^{\mathrm{c}}\setminus\mathscr{A}\big)$ for all i, as desired. ☐

Theorem 5.4. *Consider any essential* 2*-arrangement* $\mathscr{A}$ *in* S^d *or* $\mathbb{R}^d$*, let* $\mathbf{s}$ *denote the combinatorial stratification of* S^d *induced by any extension* $\mathscr{A}_e$ *of* $\mathscr{A}$*. Then the complement complex* $\mathscr{K}=\mathscr{K}(\mathscr{A}_e)$ *admits a perfect discrete Morse function.*

Proof. We distinguish two cases: (1) the case of 2-arrangements in $\mathbb{R}^d$, and (2) the case of 2-arrangements in S^d.

(1) If $\mathscr{A}$ denote any 2-arrangement of affine subspaces in $\mathbb{R}^d$, consider the radial projection ρ of $\mathbb{R}^d$ to the hemisphere $\mathcal{D}^{\mathrm{c}}$ in S^d, and the arrangement $\mathscr{A}'$ induced by $\mathscr{A}$, as defined in Definition 2.4. The complex $\mathrm{R}^*(\mathscr{K},\mathcal{D}^{\mathrm{c}})$ is a model for $\mathscr{A}$, and the discrete Morse function $X^{\mathrm{c}}_{\mathrm{R}^*(\mathscr{K},\mathcal{D}^{\mathrm{c}})}$ constructed in Lemma 5.3 provides a perfect Morse function on it, finishing the proof.

(2) Suppose instead $\mathscr{A}$ is any 2-arrangement in S^d. Let $\mathcal{D}$ denote a generic hemisphere in S^d. By Lemma 5.3, there is a discrete Morse function $\mathit{\Psi}$ on $\mathbf{s}$ whose critical faces are the subcomplex $\mathrm{R}(\mathbf{s},\mathcal{D})$ and an additional facet of $\mathbf{s}$ such that the complement matching $\mathit{\Psi}^{\mathrm{c}}_{\mathrm{R}^*[\mathcal{D}^{\mathrm{c}}]}$, restricted to $\mathrm{R}^*[\mathcal{D}^{\mathrm{c}}]$, is perfect. By Theorem 3.1(A), the matching $\mathit{\Psi}$ can be extended to a matching $\mathit{\Omega}$ on $\mathscr{K}$ such that only outwardly matched faces of dimension $\iota(d)$ with respect to the pair $(\mathbf{s},\mathrm{R}(\mathbf{s},\mathscr{A}))$ are added. Thus, $\Sigma(\Omega^{\mathrm{c}})$ is obtained from $\Sigma(\mathit{\Psi}^{\mathrm{c}}_{\mathrm{R}^*[\mathcal{D}^{\mathrm{c}}]})$ by attaching e-dimensional cells, where $e=\lceil d/2\rceil = d-\iota(d)-1$. Each of these cells can either add a generator for homology in dimension e, or delete a generator for the homology in dimension $e-1$. But if one of the cells deletes a generator, then

$$\beta_{e-1}(\Sigma(\mathit{\Psi}^{\mathrm{c}}_{\mathrm{R}^*[\mathcal{D}^{\mathrm{c}}]})) < \beta_{e-1}(\Sigma(\Omega^{\mathrm{c}})),$$

which contradicts Lemma 5.1(I). Thus, every cell attached adds a generator in homology, and in particular, since $c_i(\mathit{\Psi}^{\mathrm{c}}_{\mathrm{R}^*[\mathcal{D}^{\mathrm{c}}\cap H]}) = \beta_i\big(\mathcal{D}^{\mathrm{c}}\setminus\mathscr{A}\big)$ for all i, we have $c_i(\mathit{\Omega}^{\mathrm{c}}) = \beta_i\big(S^d\setminus\mathscr{A}\big)$ for all i, finishing the proof.

☐

6 Appendix I: Proof of Lemma 5.1

Let us recall the Goresky–MacPherson formula. For an element p of a poset P, we denote by $P_{<p}$ the poset of elements of $P(\mathscr{A})$ smaller than p. Let $\Delta(P)$ denote the order complex of a poset P. Let $\widetilde{H}_*$ resp. $\widetilde{H}^*$ denote reduced homology resp. cohomology, and let $\widetilde{\beta}_*$ denote the reduced Betti number.

Theorem 6.1 (Goresky–MacPherson [GM88, Sec. III, Thm. A], [GM88, Cor. 2.3])**.** *Let* $\mathscr{A}$ *denote an arrangement of affine subspaces in* $\mathbb{R}^d$*. Let* $\mathcal{P}(\mathscr{A})$ *denote the poset of nonempty intersections of elements of* $\mathscr{A}$*, ordered by reverse inclusion. Then, for all* i*,*

$$\widetilde{H}^i(\mathbb{R}^d\setminus\mathscr{A};\mathbb{Z}) \cong \bigoplus_{p\in\mathcal{P}} \widetilde{H}_{d-2-i-\dim p}(\Delta(\mathcal{P}_{<p}(\mathscr{A}));\mathbb{Z}).$$

Apart from the Goresky–MacPherson formula, we will make use of the following observation: We say that a subposet Q of a poset P is a *truncation* of P if every element of P not in Q is a maximal element of P. In this case, for every element q of Q, we have that $Q_{<q}=P_{<q}$, and in particular, $\Delta(Q_{<q})=\Delta(P_{<q})$.

We turn to the proof of the claims of Lemma 5.1.

Proof of Lemma 5.1. Let $\mathscr{A}$ denote a subspace arrangement in $S^d \subset \mathbb{R}^{d+1}$. The arrangement $\mathscr{A}$ gives rise to a subspace arrangement $\mathscr{A}_{\mathrm{sp}}$ in $\mathbb{R}^{d+1}$ by considering, for every element p of $\mathscr{A}$, the subspace $\mathrm{sp}(p)$ in $\mathbb{R}^{d+1}$. Furthermore, if $\mathcal{D}$ is a general position open hemisphere (with respect to $\mathscr{A}$), and π is the central projection of $\mathcal{D}$ to $\mathbb{R}^d$, then for every element p of $\mathscr{A}$, $\pi(p)$ is an affine subspace in $\mathbb{R}^d$. The collection of subspaces $\pi(p)$, $p \in \mathscr{A}$ gives a subspace arrangement $\mathscr{A}_\pi$. Notice that $\mathbb{R}^{d+1} \setminus \mathscr{A}_{\mathrm{sp}}$ is homotopy equivalent to $S^d \setminus \mathscr{A}$, and that $\mathbb{R}^d \setminus \mathscr{A}_\pi$ is homotopy equivalent to $\mathcal{D} \setminus \mathscr{A}$.

(I) $\beta_i(S^d \setminus \mathscr{A}) \geq \beta_i(\mathcal{D} \setminus \mathscr{A})$ for all i

The poset $\mathcal{P}(\mathscr{A})$ is isomorphic to $\mathcal{P}(\mathscr{A}_\pi)$, so, for all elements p of $\mathcal{P}(\mathscr{A})$, we have

$$\Delta(\mathcal{P}_{<\pi(p)}(\mathscr{A}_\pi)) \cong \Delta(\mathcal{P}_{<p}(\mathscr{A})),$$

so by Theorem 6.1,

$$\widetilde{\beta}_i(\mathbb{R}^d \setminus \mathscr{A}_\pi) = \sum_{p \in \mathcal{P}(\mathscr{A}_\pi)} \widetilde{\beta}_{d-2-i-\dim p}\big(\Delta(\mathcal{P}_{<p}(\mathscr{A}_\pi))\big) = \sum_{p \in \mathcal{P}(\mathscr{A})} \widetilde{\beta}_{d-2-i-\dim p}\big(\Delta(\mathcal{P}_{<p}(\mathscr{A}))\big)$$

for all i. Now, the poset $\mathcal{P}(\mathscr{A})$ is isomorphic to a truncation of $\mathcal{P}(\mathscr{A}_{\mathrm{sp}})$, so for all $p \in \mathcal{P}(\mathscr{A})$

$$\Delta(\mathcal{P}_{<\mathrm{sp}(p)}(\mathscr{A}_{\mathrm{sp}})) \cong \Delta(\mathcal{P}_{<p}(\mathscr{A})),$$

so by Theorem 6.1 we have that for all i

$$\widetilde{\beta}_i(\mathbb{R}^{d+1} \setminus \mathscr{A}_{\mathrm{sp}}) = \sum_{p \in \mathcal{P}(\mathscr{A}_{\mathrm{sp}})} \widetilde{\beta}_{d-1-i-\dim p}\big(\Delta(\mathcal{P}_{<p}(\mathscr{A}_{\mathrm{sp}}))\big) \geq \sum_{\substack{p \in \mathcal{P}(\mathscr{A}_{\mathrm{sp}}) \\ p \neq 0}} \widetilde{\beta}_{d-1-i-\dim p}\big(\Delta(\mathcal{P}_{<p}(\mathscr{A}_{\mathrm{sp}}))\big),$$

where

$$\sum_{\substack{p \in \mathcal{P}(\mathscr{A}_{\mathrm{sp}}) \\ p \neq 0}} \widetilde{\beta}_{d-1-i-\dim p}\big(\Delta(\mathcal{P}_{<p}(\mathscr{A}_{\mathrm{sp}}))\big) = \sum_{p \in \mathcal{P}(\mathscr{A})} \widetilde{\beta}_{d-2-i-\dim p}\big(\Delta(\mathcal{P}_{<p}(\mathscr{A}))\big) = \widetilde{\beta}_i(\mathbb{R}^d \setminus \mathscr{A}_\pi).$$

Thus

$$\widetilde{\beta}_i(S^d \setminus \mathscr{A}) = \widetilde{\beta}_i(\mathbb{R}^d \setminus \mathscr{A}_{\mathrm{sp}}) \geq \widetilde{\beta}_i(\mathbb{R}^d \setminus \mathscr{A}_\pi) = \widetilde{\beta}_i(\mathcal{D} \setminus \mathscr{A}),$$

and thus, for all i,

$$\beta_i(S^d \setminus \mathscr{A}) \geq \beta_i(\mathcal{D} \setminus \mathscr{A}).$$

(II) If H is a general position hyperplane in S^d, then $\beta_i(\mathcal{D} \setminus \mathscr{A}) \geq \beta_i((\mathcal{D} \cap H) \setminus \mathscr{A})$ for all i

If p is an element of $\mathscr{A}_\pi$ intersecting the hyperplane $\pi(H)$, let $p^{\pi(H)} := p \cap \pi(H)$ denote the corresponding subspace of $\pi(H)$. Let $\mathscr{A}_\pi^{\pi(H)}$ denote the affine arrangement obtained as the union of $p^{\pi(H)}$, $p \in \mathscr{A}_\pi$. Now, $\mathcal{P}(\mathscr{A}_\pi^{\pi(H)})$ is isomorphic to a truncation of $\mathcal{P}(\mathscr{A}_\pi)$, so for every element p of $\mathscr{A}_\pi$ intersecting $\pi(H)$, we have

$$\Delta(\mathcal{P}_{<p^{\pi(H)}}(\mathscr{A}_\pi^{\pi(H)})) \cong \Delta(\mathcal{P}_{<p}(\mathscr{A}_\pi)).$$

By this observation, and the Goresky–MacPherson formula (Theorem 6.1), we have

$$\widetilde{\beta}_i(\pi(H) \setminus \mathscr{A}_\pi) = \sum_{p \in \mathcal{P}\left(\mathscr{A}_\pi^{\pi(H)}\right)} \widetilde{\beta}_{d-2-i-\dim p}\big(\Delta(\mathcal{P}_{<p}(\mathscr{A}_\pi^{\pi(H)}))\big) = \sum_{\substack{p \in \mathcal{P}(\mathscr{A}_\pi) \\ p \cap H \neq \emptyset}} \widetilde{\beta}_{d-2-i-\dim p}\big(\Delta(\mathcal{P}_{<p}\mathscr{A}_\pi)\big).$$

and again by Theorem 6.1, we furthermore have

$$\widetilde{\beta}_i(\mathbb{R}^d \setminus \mathscr{A}_\pi) = \sum_{p \in \mathcal{P}(\mathscr{A}_\pi)} \widetilde{\beta}_{d-2-i-\dim p}\big(\Delta(\mathcal{P}_{<p}(\mathscr{A}_\pi))\big) \geq \sum_{\substack{p \in \mathcal{P}(\mathscr{A}_\pi) \\ p \cap H \neq \emptyset}} \widetilde{\beta}_{d-2-i-\dim p}\big(\Delta(\mathcal{P}_{<p}(\mathscr{A}_\pi))\big).$$

Consequently, we have

$$\widetilde{\beta}_i(\mathcal{D} \setminus \mathscr{A}) = \widetilde{\beta}_i(\mathbb{R}^d \setminus \mathscr{A}_\pi) \geq \widetilde{\beta}_i(\pi(H) \setminus \mathscr{A}_\pi) = \widetilde{\beta}_i((\mathcal{D} \cap H) \setminus \mathscr{A}),$$

and thus we obtain $\beta_i(\mathcal{D} \setminus \mathscr{A}) \geq \beta_i((\mathcal{D} \cap H) \setminus \mathscr{A})$ for all i, as desired. ☐

7 Appendix II. Minimality of c-arrangements

Recall that a *c-arrangement $\mathscr{A}$* in S^d (resp. in $\mathbb{R}^d$) is a collection $(h_i)_{i\in[n]}$ of distinct $(d-c)$-dimensional subspaces of S^d (resp. of $\mathbb{R}^d$), such that the codimension of any non-empty intersection of its elements is divisible by c. Is the complement of any c-arrangement $\mathscr{A}$ a minimal space? For $c = 2$, the answer is positive, as we saw in Corollary II. For $c \neq 2$, the answer is also positive. In fact, the complement $\mathscr{A}^c$ of any c-arrangement $\mathscr{A}$ in S^d or $\mathbb{R}^d$, $c \neq 2$, has the following properties:

- $\mathscr{A}^c$ has the homotopy type of a CW complex. (This holds for arbitrary subspace arrangements.)
- $\mathscr{A}^c$ has no torsion in homology. (This holds for arbitrary c-arrangements, cf. Goresky–MacPherson [GM88, Sec. III Thm. B].)
- Every connected component of $\mathscr{A}^c$ is simply connected. (This is easy to prove, but holds only if $c \neq 2$.)

Now, *any topological space satisfying these three properties is a minimal space*. This is proven in Hatcher [Hat02, Prp. 4C.1]; see also Anick [Ani84, Lem. 2] and Papadima–Suciu [PS02, Rm. 2.14]. As a consequence, Corollary II can be re-stated as follows:

Corollary 7.1. *The complement of any c-arrangement is a minimal space.*

In the rest of this section, we provide an analog of Theorem I for c-arrangements. First, some definitions. A *sign extension $\mathscr{A}_\sigma$* of a c-arrangement $\mathscr{A} = (h_i)_{i\in[n]}$ in S^d is any collection of hyperplanes

$$(H_{ij}) \subset S^d,\ i \in [n],\ j \in [c-1]$$

such that $\cap_{j\in[c-1]} H_{ij} = h_i$ for each i. If I is any subset of $[c-1]$, then we say that the intersection $\cap_{j\in I} H_{ij}$ *extends* h_i. A *hyperplane extension $\mathscr{A}_e$* of $\mathscr{A}$ in S^d is a sign extension of $\mathscr{A}$ together with an arbitrary collection of hyperplanes in S^d. The subdivision of S^d into convex subsets induced by $\mathscr{A}_e$ is the *stratification of S^d given by $\mathscr{A}_e$*. A stratification is *combinatorial* if it is a regular CW complex.

Let $\mathbf{s}$ be any combinatorial stratification of the sphere S^d associated to a c-arrangement $\mathscr{A}$. Let $\mathbf{s}^*$ be the dual block complex of $\mathbf{s}$. Define $\mathscr{K}(\mathscr{A}_e) = \mathrm{R}^*(\mathbf{s}^*, S^d \setminus \mathscr{A})$. The regular CW complex $\mathscr{K}(\mathscr{A}_e)$ is the *complement complex of $\mathscr{A}$ induced by $\mathscr{A}_e$*. Complement complexes of c-arrangements in $\mathbb{R}^d$ are defined by restricting the complement complex of a spherical arrangement to a general position hemisphere, cf. Definition 2.4.

Theorem 7.2. *Let $\mathscr{K}$ be the complement complex constructed from any combinatorial stratification of any essential c-arrangement $\mathscr{A}$ in S^d or $\mathbb{R}^d$. Then $\mathscr{K}$ admits a perfect discrete Morse function.*

Proof. The proof of this theorem is identical to the proof of Theorem 5.4 up to replacing 2 by c. ☐

Acknowledgements Special thanks to Bruno Benedetti, who asked me whether complement complexes admit perfect Morse functions, and who was of substantial help when writing this paper.
Thanks to Günter M. Ziegler, Alex Suciu, Mario Salvetti and Anders Björner for helpful discussions and suggestions.
Many thanks to the Institut Mittag-Leffler, Sweden, and the Hebrew University of Jerusalem, Israel, where part of the research was conducted.

References

[AB11] K. Adiprasito and B. Benedetti, *Metric geometry and collapsibility*, preprint, `arXiv:1107.5789`.

[AB12a] ______, *Smooth and discrete Morse theory*, in preparation.

[AB12b] ______, *Subdivisions, shellability and the Zeeman conjecture*, preprint, `arXiv:1202.6606`.

[Ani84] D. Anick, *Connections between Yoneda and Pontrjagin algebras*, Algebraic Topology Aarhus 1982 (Ib Madsen and Robert Oliver, eds.), Lecture Notes in Mathematics, vol. 1051, Springer Berlin / Heidelberg, 1984, 10.1007/BFb0075575, pp. 331–350.

[Ben12] B. Benedetti, *Discrete Morse theory for manifolds with boundary*, Trans. Amer. Math. Soc. **364** (2012), 6631–6670.

[BZ92] A. Björner and G. M. Ziegler, *Combinatorial stratification of complex arrangements*, J. Amer. Math. Soc. **5** (1992), 105–149.

[CS98] D. C. Cohen and A. I. Suciu, *Homology of iterated semidirect products of free groups*, J. Pure Appl. Algebra **126** (1998), 87–120.

[DP03] A. Dimca and S. Papadima, *Hypersurface complements, Milnor fibers and higher homotopy groups of arrangements*, Ann. of Math. (2) **158** (2003), 473–507.

[Fal93] M. J. Falk, *Homotopy types of line arrangements*, Invent. Math. **111** (1993), no. 1, 139–150.

[For98] R. Forman, *Morse theory for cell complexes*, Adv. Math. **134** (1998), 90–145.

[GM88] M. Goresky and R. MacPherson, *Stratified Morse Theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), vol. 14, Springer-Verlag, Berlin, 1988.

[Goo68] R. E. Goodrick, *Non-simplicially collapsible triangulations of I^n*, Proc. Cambridge Philos. Soc. **64** (1968), 31–36.

[Hat75] A. Hattori, *Topology of C^n minus a finite number of affine hyperplanes in general position*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **22** (1975), 205–219.

[Hat02] A. Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, 2002.

[HL71] H. Hamm and D. T. Lê, *Un théorème du type de Lefschetz*, C. R. Acad. Sci. Paris Sér. A-B **272** (1971), A946–A949.

[Kir95] R. Kirby, *Problems in low–dimensional topology*, available at `http://math.berkeley.edu/~kirby/problems.ps.gz`.

[Mun84] James R. Munkres, *Elements of Algebraic Topology*, Addison-Wesley Publishing Company, Menlo Park, CA, 1984.

[OT92] P. Orlik and H. Terao, *Arrangements of Hyperplanes*, Grundlehren der Mathematischen Wissenschaften, vol. 300, Springer-Verlag, Berlin, 1992.

[PS02] S. Papadima and A. I. Suciu, *Higher homotopy groups of complements of complex hyperplane arrangements*, Advances in Mathematics **165** (2002), no. 1, 71–100.

[Ran02] R. Randell, *Morse theory, Milnor fibers and minimality of hyperplane arrangements*, Proc. Amer. Math. Soc. **130** (2002), 2737–2743.

[Ryb11] G. L. Rybnikov, *On the fundamental group of the complement of a complex hyperplane arrangement*, Funktsional. Anal. i Prilozhen. **45** (2011), 71–85.

[Sal87] M. Salvetti, *Topology of the complement of real hyperplanes in $\mathbb{C}^n$*, Invent. Math. **88** (1987), 603–618.

[SS07] M. Salvetti and S. Settepanella, *Combinatorial Morse theory and minimality of hyperplane arrangements*, Geom. Topol. **11** (2007), 1733–1766.

[Yos07] M. Yoshinaga, *Hyperplane arrangements and Lefschetz's hyperplane section theorem*, Kodai Math. J. **30** (2007), 157–194.

[Zee64] E. C. Zeeman, *On the dunce hat*, Topology **2** (1964), 341–358.

[Zie93] G. M. Ziegler, *On the difference between real and complex arrangements*, Math. Z. **212** (1993), 1–11.

[ZŽ93] G. M. Ziegler and R. T. Živaljević, *Homotopy types of arrangements via diagrams of spaces*, Math. Annalen **295** (1993), 527–548.